

LECTURES ON INTEGRABLE HIERARCHIES AND VERTEX OPERATORS

A. A. VLADIMIROV

BLTP, JINR, Dubna, Moscow region 141980, Russia
alvladim@thsun1.jinr.ru

Contents

I. INTEGRABLE HIERARCHIES	
Scalar Ansatz (KP hierarchy)	2
Fermionic Fock space	7
Fermi-Bose correspondence	10
KP hierarchy via free fermions	14
II. VERTEX ALGEBRAS	
Formal distributions and locality	15
Operator product expansion	17
Vertex algebras	19
Free fermions	20
Virasoro algebra in KdV	23

Abstract

This is a slightly extended write-up of lectures given at the 2004 DIAS-TH Winter school on theoretical physics in Dubna, Russia. The exposition is intended for undergraduate students. The two parts of this course are linked together by the concept of vertex operators, which naturally arise, via Fermi-Bose correspondence, in the context of integrable hierarchies, and later play a central role in vertex algebras, under the name of the state-field map. Also, the final section illustrates another deep connection between integrable (here KdV) hierarchies and vertex (here Virasoro) algebras. Last updates and some related material can be found on my site <http://thsun1.jinr.ru/~alvladim/qft.html>

The lectures do not include any original results of the author, and are mostly compiled from well-known textbooks and introductory papers. Main references to be cited here are:

- T. Miwa, M. Jimbo and E. Date, *Solitons*, 2000.
- V. Kac, *Infinite Dimensional Lie Algebras*, 1990.
- V. Kac, *Vertex Algebras for Beginners*, 1998.
- H. Aratyn, hep-th/9503211.
- A. Matsuo and K. Nagatomo, hep-th/9706118.
- E. Witten, Comm.Math.Phys. 114 (1988) 1.

I. INTEGRABLE HIERARCHIES

Scalar Ansatz (KP hierarchy)

Probably, the most efficient way to deal with integrable nonlinear equations is the Lax representation

$$\partial_t L = [M, L]. \quad (1)$$

This readily produces a formally infinite set of conserved quantities $I_n \sim \text{tr } L^n$:

$$(1) \Rightarrow \partial_t L^n = [M, L^n] \Rightarrow \partial_t \text{tr } L^n = \text{tr}[M, L^n] = 0, \quad (2)$$

and enables us to view the t -evolution of L as a similarity transformation,

$$\partial_t T = MT, \quad \partial_t T^{-1} = -T^{-1}M \Rightarrow L(t) = TL(0)T^{-1}, \quad (3)$$

which implies isospectrality of the corresponding eigenvalue problem:

$$L(t)\psi(t) = z\psi(t) \Rightarrow \partial_t z = 0, \quad \psi(t) = T\psi(0), \quad \partial_t \psi = M\psi. \quad (4)$$

To find a Lax (or $L - M$) pair for a given equation is an extremely difficult (and generally unsolvable) task. If one, on the contrary, starts with some reasonable Ansatz for L , it appears fairly possible to discover an infinite set $\{M_n\}$ of appropriate M -operators. If, moreover, the flows given by the corresponding equations (1) commute (so that the t_n -evolutions may be considered as independent, and the t_n themselves – as a new set of coordinates on the phase space), the whole set of equations (1) with M_n is called an (integrable) hierarchy.

In practice, integrable hierarchies are highly symmetric, infinite sets of nonlinear evolution equations of the Lax type for infinitely many functions u_i of infinitely many variables t_n , $n = 1, 2, \dots$. We use the notation $x \equiv t_1$, $\partial \equiv \partial_x \equiv \partial_1$, $\partial_n \equiv \frac{\partial}{\partial t_n}$. When needed, we will specially indicate the operator nature of a derivative, e.g., $\partial_n \circ B = \partial_n B + B \partial_n$. Formal sums of the type

$$P(x, \partial) = \sum_m a_m(x) \partial^m, \quad a_m(x) = 0 \text{ for } m \gg 0 \quad (5)$$

are called pseudodifferential operators (in a variable x . Other possible arguments of a_m are treated as parameters). As usual,

$$P_+ = \sum_{m \geq 0} a_m(x) \partial^m, \quad P_- = P - P_+ = \sum_{m < 0} a_m(x) \partial^m, \quad \text{Res } P = a_{-1}(x), \quad (6)$$

so that P_+ is a proper differential part of P . Operator ∂^{-1} is defined by $\partial \partial^{-1} = \partial^{-1} \partial = 1$. For any integer m , the following holds:

$$\partial^m \circ f = \sum_{k \geq 0} \binom{m}{k} \partial^k f \partial^{m-k}. \quad (7)$$

With this rule, pseudodifferential operators form an associative algebra. This is shown straightforwardly, as well as the following useful relation:

$$\text{Res}[P, Q] = \partial(\dots). \quad (8)$$

The Kadomtsev-Petviashvili (KP) hierarchy is generated by the following (scalar) Ansatz for the Lax operator:

$$L = \partial + \sum_{i=1}^{\infty} u_i \partial^{-i}. \quad (9)$$

Unknown functions u_i may depend on all arguments t_n , or on a part of them. We use

$$B_n \doteq (L^n)_+ \equiv L_+^n \quad (B_1 = \partial, \quad B_2 = \partial^2 + 2u_1, \quad B_3 = \partial^3 + 3\partial_x u_1 + 3u_1 \partial + 3u_2, \dots) \quad (10)$$

to formulate the infinite system of equations of the KP hierarchy:

$$\partial_n L = [B_n, L] \quad \text{or} \quad [\partial_n - B_n, L] = 0 \quad (n = 1, 2, \dots) \quad (11)$$

These are nonlinear equations in u_i . Their integrability is based on the Lax form of (11) and on the following “zero-curvature” property,

$$[\partial_m - B_m, \partial_n - B_n] = \partial_n B_m - \partial_m B_n + [B_m, B_n] = 0. \quad (12)$$

From this, one easily deduces that the flows generated by different B_n commute,

$$\partial_n [B_m, L] = \partial_m [B_n, L]. \quad (13)$$

To derive (12) itself, we observe that (11) entails $\partial_n L^m = [B_n, L^m]$, that $[B_n, L^m] = -[L_-^n, L^m]$, and then do the following calculations:

$$\begin{aligned} \partial_n L^m - \partial_m L^n &= [B_n, L^m] - [B_m, L^n] = [B_n, L^m] + [L_-^m, L^n] \\ &= [B_n, B_m] + [B_n, L_-^m] + [L_-^m, L^n] = [B_n, B_m] + [L_-^m, L_-^n]. \end{aligned} \quad (14)$$

A positive part of this equality is exactly (12).

The KP equation for $u_1 = u(x, y, t)$, $y = t_2$, $t = t_3$,

$$3u_{yy} + \partial_x(-4u_t + u_{xxx} + 12uu_x) = 0, \quad (15)$$

which gave its name to the whole hierarchy, is extracted either from (12) for $m = 2, n = 3$, or from (11) with $n = 2, 3$.

The main equations (11) can be viewed as the compatibility condition of the linear system

$$L\psi = z\psi, \quad \partial_n \psi = B_n \psi. \quad (16)$$

Here $\psi(t, z) \equiv \psi(\{t_n\}, z)$ is known as Baker’s function. It admits an expansion

$$\ln \psi = \sum_{n=1}^{\infty} z^n t_n + \sum_{i=1}^{\infty} z^{-i} \psi_i(t). \quad (17)$$

To show this, we note that ∂ , and B_m in general, can be re-expanded in powers of L :

$$\partial = L + \sum_{i=1}^{\infty} \sigma_i^{(1)} L^{-i}, \quad (18)$$

$$B_m = L^m + \sum_{i=1}^{\infty} \sigma_i^{(m)} L^{-i}, \quad (19)$$

all $\sigma_i^{(m)}$ being expressible in $\sigma_i^{(1)}$. From (16) we see that $L^m\psi = z^m\psi$ and

$$\partial_m\psi = (z^m + \sum_{i=1}^{\infty} z^{-i}\sigma_i^{(m)})\psi, \quad (20)$$

which agrees with (17) if we fix

$$\partial_m\psi_n = \sigma_n^{(m)}. \quad (21)$$

Further, we observe that

$$\partial_k\sigma_n^{(m)} = \partial_m\sigma_n^{(k)} = \partial_k\partial_m\psi_n, \quad (22)$$

and, as a corollary, $\partial_k\sigma_n^{(1)} = \partial(\dots)$, i.e., $\sigma_n^{(1)}$ are conserved currents. Another set of such currents is provided by $\sigma_1^{(m)} = -\text{Res}L^m$ (due to (19)):

$$\partial_k\sigma_1^{(m)} = -\text{Res}\partial_kL^m = -\text{Res}[B_k, L^m] = \partial(\dots). \quad (23)$$

These two sets are not independent: it can be shown that

$$\sigma_1^{(n)} - n\sigma_n^{(1)} = \sum_{m=1}^{n-1} \partial_m\sigma_{n-m}^{(1)}. \quad (24)$$

Let us briefly describe the derivation. Using in $L^{m+1} = LL^m$ eq. (18) for L and (19) for L^{m+1} and L^m , and collecting only positive contributions, we come to

$$B_{m+1} = \partial \circ B_m - \sum_{i=1}^m \sigma_i^{(1)} B_{m-i} - \sigma_1^{(m)}. \quad (25)$$

Then, acting by both sides upon ψ , we get rid of B and ψ via (16) and (20), and arrive at a general recursion relation for $\sigma_i^{(m)}$:

$$\sigma_i^{(m+1)} = \partial_m\sigma_i^{(1)} + \sigma_{i+1}^{(m)} + \sigma_{m+i}^{(1)} - \sum_{j=1}^{m-1} \sigma_j^{(1)}\sigma_i^{(m-j)} + \sum_{j=1}^{i-1} \sigma_j^{(1)}\sigma_{i-j}^{(m)}. \quad (26)$$

Setting $i = n - m$ and summing up in m cancels bilinear parts and yields exactly (24).

An important corollary of (23) is that for any k, m

$$\partial_k\partial_m\psi_1 = \partial_k\sigma_1^{(m)} = \partial(\dots), \quad (27)$$

and we are prompted to make a definition

$$\psi_1(t) \doteq -\partial \ln \tau(t). \quad (28)$$

This is the first appearance of the famous τ -function. We see that

$$\sigma_1^{(m)} = \partial_m\psi_1 = -\partial_m\partial \ln \tau, \quad u_1 = -\sigma_1^{(1)} = \partial^2 \ln \tau. \quad (29)$$

Moreover, the recursion relation (24) can be explicitly resolved for $\sigma_n^{(1)}$ in terms of τ -function via the Schur polynomials $p_n(t)$, which are defined (for natural n) as follows:

$$e^{\xi(t,z)} \doteq \sum_{n=0}^{\infty} z^n p_n(t), \quad \xi(t,z) = \sum_{i=1}^{\infty} z^i t_i. \quad (30)$$

In particular, $p_0(t) = 1$, $p_1(t) = t_1$, $p_2(t) = \frac{1}{2}t_1^2 + t_2$, $p_3(t) = \frac{1}{6}t_1^3 + t_1t_2 + t_3$. The following properties are readily deduced:

$$\partial_m p_n(t) = p_{n-m}(t), \quad np_n(\tilde{t}) = \sum_{m=1}^n t_m p_{n-m}(\tilde{t}), \quad (31)$$

where tilde traditionally indicates a special scaling transformation of (infinite number of) arguments of p_n ,

$$\tilde{t} = \{t_1, \frac{t_2}{2}, \frac{t_3}{3}, \dots\}, \quad \tilde{\partial} = \{\partial_1, \frac{\partial_2}{2}, \frac{\partial_3}{3}, \dots\}, \quad (32)$$

so that

$$\exp\left(\sum_{n=1}^{\infty} \frac{t_n}{n} z^n\right) = \sum_{n=0}^{\infty} p_n(\tilde{t}) z^n. \quad (33)$$

The second equality in (31) follows from an observation that multiplying (33) by $1 + \xi(t, z)$ is equivalent to differentiating it by $z \frac{d}{dz}$.

The explicit solution of (24) mentioned above reads

$$\sigma_n^{(1)} = p_n(-\tilde{\partial}) \partial \ln \tau. \quad (34)$$

To verify it, we replace t_m in (31) by $-\partial_m$,

$$np_n(-\tilde{\partial}) = -\sum_{m=1}^{n-1} \partial_m p_{n-m}(-\tilde{\partial}) - \partial_n, \quad (35)$$

apply this to $\partial \ln \tau$, and then compare with (24).

Now we see from $\partial \psi_i = \sigma_i^{(1)}$ that, up to irrelevant x -independent terms,

$$\ln \psi(t, z) = \sum_{n=1}^{\infty} z^n t_n + \sum_{i=1}^{\infty} z^{-i} p_i(-\tilde{\partial}) \ln \tau = \xi(t, z) + \ln \tau(\{t_m - \frac{1}{mz^m}\}) - \ln \tau(t) \quad (36)$$

because

$$f(t_1 + \frac{1}{z}, t_2 + \frac{1}{2z^2}, \dots) = \exp\left(\sum_{n=1}^{\infty} \frac{\partial_n}{nz^n}\right) f(t) = \sum_{n=0}^{\infty} z^{-n} p_n(\tilde{\partial}) f(t). \quad (37)$$

Thus, we are able to express the Baker function ψ in terms of τ :

$$\psi(t, z) = e^{\xi(t, z)} \frac{\tau(\{t_m - \frac{1}{mz^m}\})}{\tau(t)}. \quad (38)$$

Using again (36) and (35), we can also ‘invert’ this relation and express derivatives of τ in terms of ψ :

$$\partial_n \ln \tau(t) = -\text{Res}_z z^n \left(\sum_{m=1}^{\infty} z^{-m-1} \partial_m - \partial_z \right) \ln w(t, z), \quad (39)$$

where Res_z means the coefficient of z^{-1} , and

$$\psi(t, z) = w(t, z) e^{\xi(t, z)}, \quad w(t, z) = 1 + \sum_{i=1}^{\infty} w_i(t) z^{-i}. \quad (40)$$

It should be noted here that a class of valid objects to act upon with pseudodifferential operators according to the rules $\partial^m e^{\xi(t,z)} = z^m e^{\xi(t,z)}$ and (7) is given by $(\sum_{n \leq N} z^n b_n(t)) e^{\xi(t,z)}$.

Due to (40), it seems now natural to try the following representation for ψ :

$$\psi(t, z) = W e^{\xi(t,z)}, \quad (41)$$

where the *dressing operator* W is of the form

$$W = 1 + \sum_{i=1}^{\infty} w_i(t) \partial^{-i}. \quad (42)$$

Owing to $\partial_n e^{\xi(t,z)} = \partial^n e^{\xi(t,z)} = z^n e^{\xi(t,z)}$, we see from (17), (36), (41) and (42), that w_i , ψ_i and derivatives of τ are related by

$$\sum_{i=1}^{\infty} z^{-i} \psi_i = \ln(1 + \sum_{i=1}^{\infty} z^{-i} w_i) = \sum_{i=1}^{\infty} z^{-i} p_i(-\tilde{\partial}) \ln \tau. \quad (43)$$

To guarantee (16), operator W should satisfy the following “dressing” relations:

$$L = W \partial \circ W^{-1}, \quad \partial_n - B_n = W(\partial_n - \partial^n) \circ W^{-1}. \quad (44)$$

The last of them reveals the nature of commutativity of the operators $\partial_n - B_n$: they are nothing more than dressed counterparts of the (trivially commuting) operators $\partial_n - \partial^n$. Eqs. (44) can also be rewritten in the form involving W only,

$$\partial_n W = -(W \partial^n \circ W^{-1})_- W. \quad (45)$$

It looks quite feasible that an analogous equation for the τ -function in closed form also exists. This is really so: one can obtain a fundamental *bilinear identity*

$$\text{Res}_z \tau(\{t_m - \frac{1}{mz^m}\}) \tau(\{t'_m + \frac{1}{mz^m}\}) e^{\xi(t-t',z)} = 0. \quad (46)$$

To derive (46), it is useful to introduce a so-called adjoint Baker function

$$\psi^*(t, z) \doteq (W^*)^{-1} e^{-\xi(t,z)} = w^*(t, z) e^{-\xi(t,z)}. \quad (47)$$

In W^* , the asterisk (formal conjugation) changes sign of ∂ and the order of all operators,

$$W^* = 1 + \sum_{i=1}^{\infty} (-\partial)^{-i} \circ w_i(t), \quad (W^*)^{-1} = 1 + \sum_{i=1}^{\infty} w_i^*(t) (-\partial)^{-i}, \quad (48)$$

whereas w^* is a mere notation. In analogy with the above reasoning, one can derive

$$w^*(t, z) = 1 + \sum_{i=1}^{\infty} w_i^*(t) z^{-i} = \frac{\tau(\{t_m + \frac{1}{mz^m}\})}{\tau(t)}, \quad (49)$$

and thus rewrite the bilinear identity as follows:

$$\text{Res}_z \psi(t, z) \psi^*(t', z) = 0. \quad (50)$$

Bearing in mind the Taylor expansion in $t - t'$, it is evidently sufficient to show that

$$\text{Res}_z \partial^m \psi(t, z) \psi^*(t, z) = 0, \quad m \geq 0, \quad (51)$$

because other derivatives ($\partial_{n>1}$) can be treated via (16). First, for any two pseudodifferential operators $P = \sum a_i \partial^i$ and $Q = \sum b_i \partial^i$ one proves by direct calculation that

$$\text{Res}_z(Pe^{xz})(Qe^{-xz}) = \text{Res}PQ^* = \sum_{i+j=-1} (-)^j a_i b_j \quad (52)$$

(recall that $\text{Res} = \text{Res}_\partial$). Now (51) follows:

$$\begin{aligned} \text{Res}_z \partial^m \psi(t, z) \psi^*(t, z) &= \text{Res}_z \partial^m (W e^{\xi(t, z)}) (W^*)^{-1} e^{-\xi(t, z)} \\ &= \text{Res}_z (\partial^m \circ W e^{xz}) (W^*)^{-1} e^{-xz} = \text{Res} \partial^m \circ W W^{-1} = \text{Res} \partial^m = 0. \end{aligned} \quad (53)$$

Conversely, (16) can be derived from (50). In brief: given ψ , we reconstruct W by (42), define L by (44), show that $L\psi = z\psi$ and $(\partial_n - B_n)\psi = (\partial_n - L^n + L_-^n)\psi = e^\xi \mathcal{O}(z^{-1})$. But, due to $\text{Res}_z(\partial_n - B_n)\psi(t, z) \cdot \psi^*(t', z) = 0$, this implies $(\partial_n - B_n)\psi = 0$ (it is shown by setting $t' = t$ after differentiating by t' 0, 1, ... times). Thus, bilinear identities (46), (50) turn out to be equivalent to entire KP hierarchy (11). We shall see below that bilinear relations of this kind are naturally interpreted within the fermionic representation of the τ and Baker functions.

Fermionic Fock space

Remarkably, central objects of the KP hierarchy considered above (Baker's and τ -functions, bilinear equations, and some others) can be represented, and efficiently dealt with, in the framework of free fermionic picture, via so-called Fermi-Bose correspondence. This also reveals the underlying group theory structure of integrable hierarchies.

We begin with description of an appropriate fermionic Fock space. Consider an algebra of the free fermion operators,

$$[\psi_m, \psi_n]_+ = [\bar{\psi}_m, \bar{\psi}_n]_+ = 0, \quad [\psi_m, \bar{\psi}_n]_+ = \delta_{mn}, \quad (54)$$

with arbitrary integer indices m, n . We shall call ψ_n ($\equiv \psi_n^+$) the wedging (or creation) and $\bar{\psi}_n$ ($\equiv \psi_n^-$) the contraction (or annihilation) operators. Of course, these are generators of the Clifford algebra:

$$e_n = \psi_n + \bar{\psi}_n, \quad e_{\bar{n}} = i(\psi_n - \bar{\psi}_n), \quad [e_\alpha, e_\beta]_+ = 2\delta_{\alpha\beta}. \quad (55)$$

Let $|\Omega\rangle$ (fake vacuum) be a state annihilated by each $\bar{\psi}_n$, and $|0\rangle$ (physical vacuum) the following formal object:

$$|0\rangle = \psi_{-1}\psi_{-2}\dots|\Omega\rangle, \quad \bar{\psi}_n|0\rangle = 0. \quad (56)$$

Then

$$\psi_{n<0}|0\rangle = \bar{\psi}_{n\geq 0}|0\rangle = 0, \quad (57)$$

and we can interpret the free fermionic operators as follows:

- $\psi_{n\geq 0}$ – creation of a particle
- $\bar{\psi}_{n\geq 0}$ – annihilation of a particle
- $\bar{\psi}_{n<0}$ – creation of a hole
- $\psi_{n<0}$ – annihilation of a hole

It is also useful to introduce, for any integer m , the vectors

$$|m\rangle = \psi_{m-1}\psi_{m-2}\dots|\Omega\rangle, \quad (58)$$

so that

$$\psi_{n<m}|m\rangle = \bar{\psi}_{n\geq m}|m\rangle = 0. \quad (59)$$

The fermionic Fock space $\mathbf{F}$ is spanned by vectors $|u\rangle$ which are obtained by application of a finite number of $\psi_m, \bar{\psi}_n$ operators to $|0\rangle$. So, $|\Omega\rangle$ is not in $\mathbf{F}$. We also introduce the notion of *charge* to characterize the total number of operators ψ used (in comparison with the vacuum $|0\rangle$). Thus, the charges of $\psi_n, \bar{\psi}_n$ and $|m\rangle$ are $+1, -1$ and m , respectively.

Let us now define the fermion fields as the following formal expansions:

$$\begin{aligned} \psi(z) &= \sum_n z^n \psi_n = \psi_-(z) + \psi_+(z), & \psi_-(z) &= \sum_{n<0} z^n \psi_n, & \psi_+(z) &= \sum_{n\geq 0} z^n \psi_n, & (60) \\ \bar{\psi}(z) &= \sum_n z^{-n-1} \bar{\psi}_n = \bar{\psi}_+(z) + \bar{\psi}_-(z), & \bar{\psi}_+(z) &= \sum_{n<0} z^{-n-1} \bar{\psi}_n, & \bar{\psi}_-(z) &= \sum_{n\geq 0} z^{-n-1} \bar{\psi}_n, \\ [\psi(z), \bar{\psi}(w)]_+ &= \sum_n w^n z^{-n-1} = \delta(z-w), & & & & & (61) \end{aligned}$$

see (125) for a discussion of this definition of δ -function. We see that the $(-)$ part of a field corresponds to negative, whereas the $(+)$ part to non-negative powers of z , but a more deep mnemonic rule is that $+$ corresponds to creation and $-$ to annihilation:

$$\psi(z)|0\rangle = \psi_+(z)|0\rangle, \quad \bar{\psi}(z)|0\rangle = \bar{\psi}_+(z)|0\rangle. \quad (62)$$

The duality (bra vs ket) properties of $\mathbf{F}$ are self-evident, if we accept ψ_n and $\bar{\psi}_n$ to be mutually Hermitean conjugate. We denote $\langle 0|\dots|0\rangle$ by $\langle \dots \rangle$ and use $\langle m|m\rangle = 1$. Obviously,

$$\langle \bar{\psi}_m \psi_n \rangle = -\langle \psi_n \bar{\psi}_m \rangle + \delta_{mn} = \delta_{mn} \theta(n), \quad \theta(n) = \begin{cases} 1 & (n \geq 0) \\ 0 & (n < 0) \end{cases} \quad (63)$$

Another useful example of a matrix element is (see (124))

$$\langle \psi(z) \bar{\psi}(w) \rangle = \langle \bar{\psi}(z) \psi(w) \rangle = \sum_{n \geq 0} w^n z^{-n-1} = (z-w)_w^{-1}. \quad (64)$$

The normal product is defined as usual: annihilation (with respect to the physical vacuum $|0\rangle$) operators should not stand to the left of the creation ones. For φ and χ linear in $\psi, \bar{\psi}$, it is

$$:\varphi\chi: \doteq \varphi_+\chi - \chi\varphi_- = \varphi\chi - \langle \varphi\chi \rangle = \varphi\chi - \langle \varphi_-\chi_+ \rangle. \quad (65)$$

In particular,

$$:\bar{\psi}_m \psi_n: = -:\psi_n \bar{\psi}_m: = \bar{\psi}_m \psi_n - \delta_{mn} \theta(n). \quad (66)$$

Now we introduce a bilinear current

$$J(z) = :\psi(z) \bar{\psi}(z): = \sum_n z^{-n-1} J_n, \quad J_n = \sum_m :\psi_m \bar{\psi}_{n+m}: \quad (67)$$

In fact, the normal ordering is effective only for J_0 :

$$J_0 = \sum_{m \geq 0} \psi_m \bar{\psi}_m - \sum_{m < 0} \bar{\psi}_m \psi_m. \quad (68)$$

The following holds:

$$(J_n)^+ = J_{-n}, \quad J_0|m\rangle = m|m\rangle, \quad J_{n>0}|m\rangle = 0, \quad (69)$$

so that J_0 measures the charge. The last equality in (69) is not surprising because $J_{n>0}$ effectively shifts particles by n positions down, which is clearly impossible in a ‘filled’ state $|m\rangle$. Moreover, for any Fock vector $|u\rangle$ and sufficiently large n ,

$$J_n|u\rangle = 0 \quad (n \gg 0). \quad (70)$$

Further,

$$[J_n, \psi_m] = \psi_{m-n}, \quad [J_n, \bar{\psi}_m] = -\bar{\psi}_{m+n}, \quad (71)$$

$$[J_n, \psi(z)] = z^n \psi(z), \quad [J_n, \bar{\psi}(z)] = -z^n \bar{\psi}(z). \quad (72)$$

At last, the commutator of the current components has a typical bosonic form:

$$\begin{aligned} [J_m, J_n] &= [J_m, \sum_{k \geq 0} \psi_k \bar{\psi}_{n+k} - \sum_{k < 0} \bar{\psi}_{n+k} \psi_k] \\ &= \sum_{k \geq 0} (\psi_{k-m} \bar{\psi}_{n+k} - \psi_k \bar{\psi}_{n+m+k}) - \sum_{k < 0} (-\bar{\psi}_{n+m+k} \psi_k + \bar{\psi}_{n+k} \psi_{k-m}) \\ &= \sum_{l=-m}^{-1} (\psi_l \bar{\psi}_{l+n+m} + \bar{\psi}_{l+n+m} \psi_l) = m \delta_{m,-n}. \end{aligned} \quad (73)$$

The current $J(z)$ contains infinite summation. To work with such operators in the Fock space, we need some special techniques. By definition, any Fock vector is *finite* in the following sense: it differs from $|0\rangle$ in a finite number of positions (filled or empty), or, in other words, the maximum $|n|$ of a position whose status differs from that in $|0\rangle$, is finite. When treating Fock vectors as infinite columns (the occupation number representation), it is clear that a finite vector has only finite number of nonzero components. We are interested in the operators which send finite vectors to finite. These are described by so-called finite matrices, which obey $A_{ij} = 0$ for $|i - j| \gg 0$ (so a finite matrix may actually have infinite number of nonzero elements). Multiplication (and commutation) operations respect finiteness.

Now consider a class of operators in F of the form

$$X_A = \sum_{mn} A_{mn} : \psi_m \bar{\psi}_n : \quad (74)$$

with A finite. Such operators carry zero charge, and send Fock vectors to Fock vectors: normal ordering cuts off the regions $n \gg 0$ and $m \ll 0$, and finiteness of A completes the job. For instance,

$$J_n = X_{A_n}, \quad (A_n)_{mk} = \delta_{k,n+m}. \quad (75)$$

One easily deduces

$$[X_A, \psi_n] = A_{mn} \psi_m, \quad [X_A, \bar{\psi}_n] = -A_{nm} \bar{\psi}_m, \quad (76)$$

and

$$[X_A, X_B] = X_{[A,B]} + \omega(A, B) \quad (77)$$

where the 2-cocycle

$$\omega(A, B) = -\omega(B, A) = \sum_{ij} A_{ij} B_{ji} (\theta(j) - \theta(i)) \quad (78)$$

is well defined for finite A and B . For example,

$$\omega(A_m, A_n) = m \delta_{m, -n}. \quad (79)$$

Thus we come to a centrally extended Lie algebra

$$\mathfrak{gl}_\infty = \{X_A\} \oplus \mathbb{C}. \quad (80)$$

Really, the basic commutator

$$[\psi_m \bar{\psi}_n, \psi_p \bar{\psi}_q] = \delta_{pn} \psi_m \bar{\psi}_q - \delta_{qm} \psi_p \bar{\psi}_n \quad (81)$$

copies the commutator of the (gl)-generators $(E_{mn})_{ij} = \delta_{im} \delta_{jn}$.

The corresponding Lie group $\mathbf{G}$ (a subgroup of the Clifford group) consists of invertible elements g of the Clifford algebra obeying

$$gVg^{-1} \subset V, \quad g\bar{V}g^{-1} \subset \bar{V}, \quad (82)$$

with V being a linear space spanned by ψ_n and $\bar{V}$ by $\bar{\psi}_n$. Its connected subgroup containing unity is formed by the elements of the type $\exp(X_A)$.

Consider as an example the Clifford algebra generated by $[a, a^\dagger]_+ = 1$, $aa = a^\dagger a^\dagger = 0$. Its basis is $\{1, a, a^\dagger, a^\dagger a\}$, zero-charge elements are 1 and $a^\dagger a$, whereas V and $\bar{V}$ are one-dimensional spaces along a and $a^\dagger$, respectively. The elements of the group of our interest are

$$\alpha(1 + \beta a^\dagger a) \quad \text{with} \quad \alpha \neq 0, \beta \neq -1. \quad (83)$$

Namely,

$$(1 + \beta a^\dagger a)^{-1} = (1 - \frac{\beta}{1 + \beta} a^\dagger a). \quad (84)$$

For $\beta > -1$ elements (83) admit exponential form due to

$$e^{\gamma a^\dagger a} = 1 + (e^\gamma - 1) a^\dagger a, \quad (85)$$

otherwise they don't, being in the second connected component (without unity).

We claim that any element of the orbit $\mathbf{G}|0\rangle$ obeys bilinear relations specific to the τ -function of the KP hierarchy, and thus is nothing more than a fermionic appearance of this function. To prove this (and other) statements about the properties of τ and Baker functions, and to relate their fermionic representations with more conventional ones, we need to discuss the Fermi-Bose correspondence.

Fermi-Bose correspondence

There is an isomorphism between the fermionic Fock space $\mathbf{F}$ and the (bosonic Fock) space $\mathbf{B}$ of polynomials in t_n, y, y^{-1} . Let us define

$$H(t) = \sum_{n>0} t_n J_n. \quad (86)$$

It is easily seen that

$$H(t)|m\rangle = 0, \quad [H(t), J_n] = (\theta(n) - 1)nt_{-n}. \quad (87)$$

Further useful properties of $H(t)$ are formulated in terms of $\xi(t, z)$ (30):

$$[H(t), \psi(z)] = \xi(t, z)\psi(z), \quad [H(t), \bar{\psi}(z)] = -\xi(t, z)\bar{\psi}(z), \quad (88)$$

and, consequently,

$$e^{H(t)}\psi(z)e^{-H(t)} = e^{\xi(t, z)}\psi(z), \quad e^{H(t)}\bar{\psi}(z)e^{-H(t)} = e^{-\xi(t, z)}\bar{\psi}(z). \quad (89)$$

With the use of Schur polynomials (30) one can also write

$$e^{H(t)}\psi_n e^{-H(t)} = \sum_{m=0}^{\infty} p_m(t)\psi_{n-m}, \quad (90)$$

$$e^{H(t)}\bar{\psi}_n e^{-H(t)} = \sum_{m=0}^{\infty} p_m(-t)\bar{\psi}_{n+m}. \quad (91)$$

Now, the *bosonization map* is

$$B(|u\rangle) \doteq \sum_m y^m \langle m | e^{H(t)} | u \rangle. \quad (92)$$

For any Fock vector $|u\rangle$, $B(|u\rangle)$ is a polynomial in t_n, y, y^{-1} , which is identical zero only for $|u\rangle = 0$. Clearly, $B(|m\rangle) = y^m$. Also, if $|u\rangle$ carries a definite charge m , only the y^m -term of the sum in (92) survives. Furthermore, if $|u\rangle$ is a monomial in $\psi, \bar{\psi}$ acting upon $|0\rangle$, then its Bose-counterpart can be explicitly written in terms of the Schur polynomials. Really, such a monomial $|\varphi\rangle$ is fully characterized by its charge (say, m) and a *partition* λ_φ which is a finite non-increasing sequence of natural numbers $\{\lambda_1, \lambda_2, \dots\}$, indicating the differences in positions of particles in $|\varphi\rangle$ and $|m\rangle$, beginning from the top. A general formula

$$B(|\varphi\rangle) = y^m \langle m | e^{H(t)} | \varphi \rangle = y^m p_{\lambda_\varphi}(t) \quad (93)$$

results from repeated use of (90). The Schur polynomials $p_\lambda(t)$ indexed by partitions are defined through elementary Schur polynomials (30) as follows (we assume $p_{n<0}(t) = 0$):

$$p_\lambda(t) = \det(p_{\lambda_i + j - i}(t)), \quad 1 \leq i, j \leq \sum_k \lambda_k. \quad (94)$$

For example, the state $|\varphi\rangle = \psi_m \psi_{m-1} |m-2\rangle$ has $\lambda_\varphi = \{1, 1\}$ and

$$p_{\lambda_\varphi}(t) = \det \begin{pmatrix} p_1(t) & p_2(t) \\ p_0(t) & p_1(t) \end{pmatrix} = p_1^2(t) - p_0(t)p_2(t). \quad (95)$$

Let us now derive, for later use, a formula

$$\langle m+1 | \psi(z) | \varphi \rangle = z^m p_{\lambda_\varphi}(\{-\frac{1}{nz^n}\}) = z^m \langle m | e^{-H(\{-\frac{1}{nz^n}\})} | \varphi \rangle. \quad (96)$$

To produce a nonzero contribution, a monomial $|\varphi\rangle$ (of charge m) should be equal to $|m\rangle$, or display exactly one vacancy (hole) in its ‘body’ to be filled in by some ψ_n from $\psi(z)$. Namely ($N = 0, 1, 2, \dots$)

$$|\varphi\rangle = \psi_m \psi_{m-1} \dots \psi_{m-N+1} |m-N\rangle \quad \Rightarrow \quad \langle m+1 | \psi(z) | \varphi \rangle = (-)^N z^{m-N}. \quad (97)$$

In other words, a partition λ_φ may contain only 1's. To compute the corresponding determinant in (94), observe that

$$\sum_{n=0}^{\infty} p_n(\{-\frac{1}{mz^m}\}) w^n = \exp(-\sum_{n=1}^{\infty} \frac{w^n}{nz^n}) = \exp \ln(1 - \frac{w}{z}) = 1 - \frac{w}{z}, \quad (98)$$

so

$$p_1(\{-\frac{1}{mz^m}\}) = -\frac{1}{z}, \quad p_n(\{-\frac{1}{mz^m}\}) = 0 \quad \text{for } n \geq 2. \quad (99)$$

Now the determinants in p_{λ_φ} in eq. (96) are easily evaluated. They either agree with (97), for appropriate $|\varphi\rangle$, or vanish due to (99), in both cases verifying (96). Due to $|\varphi\rangle$ being arbitrary monomial we, in fact, have proved the following useful formula:

$$\langle m|\psi(z) = z^{m-1} \langle m-1|e^{-H(\{\frac{1}{nz^n}\})}. \quad (100)$$

Analogously, one derives

$$\langle m|\bar{\psi}(z) = z^{-m-1} \langle m+1|e^{H(\{\frac{1}{nz^n}\})}. \quad (101)$$

Let us now proceed with bosonization of various fermionic operators. From (87), due to $\langle m|J_0 = m\langle m|$, $\langle m|J_{n<0} = 0$, we obtain

$$B(J_0|u) = y\partial_y B(|u) \quad (102)$$

(hence, q^{J_0} bosonizes to the operation $y \rightarrow qy$, i.e., $B(q^{J_0}|u)(t, y) = B(|u)(t, qy)$), and

$$B(J_n|u) = \partial_n B(|u), \quad B(J_{-n}|u) = nt_n B(|u) \quad (n > 0). \quad (103)$$

One concludes that components of the current $J(z)$ are represented in $\mathbf{B}$ as follows ($n > 0$):

$$J_0 \sim y\partial_y = \partial_{\ln y}, \quad J_n \sim \partial_n, \quad J_{-n} \sim nt_n, \quad (104)$$

which completely agrees with their commutation rules (73).

It appears useful to introduce in $\mathbf{F}$ an operator Y which augments charge by one and bosonizes to multiplication by y : $B(Y|u) = yB(|u)$. This is achieved via the definition

$$Y\psi_n = \psi_{n+1}Y, \quad Y^+ = Y^{-1} \quad (105)$$

that leads to the following relations:

$$Y\bar{\psi}_n = \bar{\psi}_{n+1}Y, \quad Y|m\rangle = |m+1\rangle, \quad \langle m|Y = \langle m-1|, \quad (106)$$

$$[J_n, Y] = 0 \quad (n \neq 0), \quad [J_0, Y] = Y, \quad q^{J_0}Y = qYq^{J_0}. \quad (107)$$

Taken in the form $[J_0, \ln Y] \sim [\partial_{\ln y}, \ln y] = 1$, this suggests to interpret $\ln Y$ as a creation and J_0 as an annihilation operator.

Now we are in a position to bosonize fermion fields.

$$\begin{aligned} B(\psi(z)|u) &= \sum_m y^m \langle m|e^{H(t)}\psi(z)|u\rangle = e^{\xi(t,z)} \sum_m y^m \langle m|\psi(z)e^{H(t)}|u\rangle \\ &= e^{\xi(t,z)} \sum_m y^m z^{m-1} \langle m-1|e^{H(\{t_n - \frac{1}{nz^n}\})}|u\rangle = e^{\xi(t,z)} e^{-\xi(\bar{\partial}, z^{-1})} \sum_m y^m z^{m-1} \langle m-1|e^{H(t)}|u\rangle \\ &= e^{\xi(t,z)} e^{-\xi(\bar{\partial}, z^{-1})} y \sum_m (zy)^m \langle m|e^{H(t)}|u\rangle = e^{\xi(t,z)} e^{-\xi(\bar{\partial}, z^{-1})} y z^{y\partial_y} B(|u). \end{aligned} \quad (108)$$

Performing a similar calculation for $\bar{\psi}$ we conclude that

$$\psi^\pm(z) \sim e^{\pm\xi(t,z)} e^{\mp\xi(\bar{\partial}, z^{-1})} y^{\pm 1} z^{\pm y \partial_y} = e^{\pm \sum_{n>0} z^n t_n} e^{\mp \sum_{n>0} \frac{\partial_n}{n z^n}} y^{\pm 1} z^{\pm y \partial_y}. \quad (109)$$

In view of the bosonized form of J_n , this implies

$$\psi^\pm(z) = \exp\left(\pm \sum_{m=1}^{\infty} \frac{z^m}{m} J_{-m}\right) \exp\left(\mp \sum_{n=1}^{\infty} \frac{J_n}{n z^n}\right) Y^{\pm 1} z^{\pm J_0} \quad (110)$$

and may be considered as properly ordered bosonic counterpart of $:e^{\pm\varphi(z)}:$ with

$$\varphi(z) = \sum_{n \neq 0} \frac{J_n}{-n} z^{-n} + J_0 \ln z + \ln Y, \quad \frac{d}{dz} \varphi(z) = J(z). \quad (111)$$

Clearly, in our derivation (108) of the bosonization formulas (109),(110) for the fermion fields, the relations (100),(101) play the crucial role. It should be noted that the inverted calculation, i.e., deducing (100),(101) from (110), is even simpler. So, in principle, all the construction could be started with deriving (110) from the ‘first principles’: namely, from the commutators (72). For example, the fermionic counterpart of $e^{-\xi(t,z)} \psi(z) e^{\xi(\bar{\partial}, z^{-1})}$ proves to commute with J_n for $n \neq 0$ and is thus equal to $Y f(z, J_0)$ (because in $\mathcal{B}$ this means commutativity with t_n and ∂_n , and, as a result, independence of both). A function f is found (say, using $\langle m+1 | \psi(z) | m \rangle = z^m$) to be z^{J_0} , which justifies (110).

Exponential operators like those encountered in (109) and (110),

$$V^\alpha(z) = \exp\left(\alpha \sum_{n>0} z^n t_n\right) \exp\left(-\alpha \sum_{n>0} \frac{\partial_n}{n z^n}\right) y^\alpha z^{\alpha y \partial_y} \quad (112)$$

as well as their F-counterparts, are known as *vertex operators*. Of course, $\psi^\pm(z) \sim V^{\pm 1}(z)$. From (107) and the well-known identity

$$[a, b] = \lambda \in \mathbb{C} \quad \implies \quad e^a e^b = e^\lambda e^b e^a \quad (113)$$

we find $z^{\alpha y \partial_y} y^\beta = z^{\alpha \beta} y^\beta z^{\alpha y \partial_y}$,

$$\begin{aligned} & \left[-\alpha \sum_{m=1}^{\infty} \frac{\partial_m}{m z^m}, \beta \sum_{n=1}^{\infty} w^n t_n \right] = -\alpha \beta \sum_{n=1}^{\infty} \frac{w^n}{n z^n} = \alpha \beta \ln\left(1 - \frac{w}{z}\right), \\ & \exp\left(-\alpha \sum_{m=1}^{\infty} \frac{\partial_m}{m z^m}\right) \exp\left(\beta \sum_{n=1}^{\infty} w^n t_n\right) = \frac{(z-w)_w^{\alpha \beta}}{z^{\alpha \beta}} \exp\left(\beta \sum_{n=1}^{\infty} w^n t_n\right) \exp\left(-\alpha \sum_{m=1}^{\infty} \frac{\partial_m}{m z^m}\right), \end{aligned}$$

and finally obtain

$$\begin{aligned} V^\alpha(z) V^\beta(w) &= (z-w)_w^{\alpha \beta} :V^\alpha(z) V^\beta(w): \\ &= (z-w)_w^{\alpha \beta} e^{\alpha \xi(t,z) + \beta \xi(t,w)} e^{-\alpha \xi(\bar{\partial}, z^{-1}) - \beta \xi(\bar{\partial}, w^{-1})} y^{\alpha + \beta} (z^\alpha w^\beta)^{y \partial_y}. \end{aligned} \quad (114)$$

Let us consider some particular cases. Using (114) for $\alpha = 1, \beta = -1$ together with an identity $(z-w)_w^{-1} - (z-w)_z^{-1} = \delta(z-w)$ (see (131)), one easily checks (61):

$$[\psi(z), \bar{\psi}(w)]_+ \sim e^{\xi(t,z) - \xi(t,w)} e^{-\xi(\bar{\partial}, z^{-1}) + \xi(\bar{\partial}, w^{-1})} \left(\frac{z}{w}\right)^{y \partial_y} ((z-w)_w^{-1} + (w-z)_z^{-1}) = \delta(z-w).$$

Analogously, one can verify that, for example,

$$\psi(z) \psi(w) \sim (z-w) e^{\xi(t,z) + \xi(t,w)} e^{-\xi(\bar{\partial}, z^{-1}) - \xi(\bar{\partial}, w^{-1})} y^2 (zw)^{y \partial_y} \quad (115)$$

is manifestly antisymmetric, as it should be due to fermionic nature of ψ .

KP hierarchy via free fermions

Now we are to justify our claim that any $g \in \mathbf{G}$ is the fermionic counterpart of a τ -function, i.e., of some solution of the (bilinear identities of) KP hierarchy. Namely, we offer the following bosonized expressions as candidates for the role of τ and Baker functions:

$$\tau(t) \equiv \tau(t; g) = \langle 0 | e^{H(t)} g | 0 \rangle, \quad (116)$$

$$\psi(t, z) = \frac{\langle 1 | e^{H(t)} \psi(z) g | 0 \rangle}{\tau(t; g)}, \quad (117)$$

$$\psi^*(t, z) = \frac{\langle -1 | e^{H(t)} \bar{\psi}(z) g | 0 \rangle}{\tau(t; g)}. \quad (118)$$

To begin with, we check the relation (38) using (89) and (100):

$$\begin{aligned} \langle 1 | e^{H(t)} \psi(z) g | 0 \rangle &= e^{\xi(t, z)} \langle 1 | \psi(z) e^{H(t)} g | 0 \rangle = e^{\xi(t, z)} \langle 0 | e^{H(t) - H(\{\frac{1}{nz^n}\})} g | 0 \rangle \\ &= e^{\xi(t, z)} \langle 0 | e^{H(\{t_n - \frac{1}{nz^n}\})} g | 0 \rangle = e^{\xi(t, z)} \tau(\{t_n - \frac{1}{nz^n}\}). \end{aligned} \quad (119)$$

Similarly, using (101) serves to confirm (49).

Since the proper relations between the τ and Baker functions are thus verified, it suffices to check the bilinear identity in the form (50):

$$\begin{aligned} \text{Res}_z \langle 1 | e^{H(t)} \psi(z) g | 0 \rangle \langle -1 | e^{H(t')} \bar{\psi}(z) g | 0 \rangle &= \sum_n \langle 1 | e^{H(t)} \psi_n g | 0 \rangle \langle -1 | e^{H(t')} \bar{\psi}_n g | 0 \rangle \\ &= \left(\langle 1 | e^{H(t)} \otimes \langle -1 | e^{H(t')} \right) \sum_n \psi_n g | 0 \rangle \otimes \bar{\psi}_n g | 0 \rangle. \end{aligned} \quad (120)$$

But any element $|u\rangle = g|0\rangle$ of the orbit $\mathbf{G}|0\rangle$ obeys a bilinear relation

$$\text{Res}_z \psi(z) |u\rangle \otimes \bar{\psi}(z) |u\rangle = \sum_n \psi_n |u\rangle \otimes \bar{\psi}_n |u\rangle = 0 \quad (121)$$

which is readily shown in the infinitesimal form $g = e^{X_A} \approx 1 + X_A$ with the help of (76):

$$\begin{aligned} \sum_n \psi_n g | 0 \rangle \otimes \bar{\psi}_n g | 0 \rangle &\approx \sum_n (\psi_n | 0 \rangle \otimes \bar{\psi}_n | 0 \rangle + \psi_n X_A | 0 \rangle \otimes \bar{\psi}_n | 0 \rangle + \psi_n | 0 \rangle \otimes \bar{\psi}_n X_A | 0 \rangle) \\ &= (\text{id} \otimes \text{id} + X_A \otimes \text{id} + \text{id} \otimes X_A) \sum_n \psi_n | 0 \rangle \otimes \bar{\psi}_n | 0 \rangle = 0 \end{aligned} \quad (122)$$

because for any n one of $\psi_n, \bar{\psi}_n$ must be an annihilation operator. The bilinear identity (50) is thus proved. Its another version (46), in terms of the τ -function, follows from (121) through the bosonization of fermion fields.

Of course, the orbit $\mathbf{G}|0\rangle$ provides a great supply of solutions of bilinear identities (hence, of the KP hierarchy). The simplest one is $\tau = 1$ which corresponds to the vacuum state itself. Further, any zero-charged monomial $|\varphi\rangle$ (see (93)) lies in (a completion of) the orbit, therefore all Schur polynomials $p_{\lambda_\varphi}(t)$ are solutions. Another important class of (N -soliton) solutions is produced by the (repeated) action of exponents of vertex operators, related to quadratic combinations of the Fermi fields, upon 1.

II. VERTEX ALGEBRAS

Formal distributions and locality

Vertex algebras encode mathematical content of 2-dimensional conformal quantum field theory (CFT₂). Roughly speaking, chiral quantum fields depending on a single complex variable z are replaced by operator-valued formal power series in an abstract variable z . We begin with a brief exposition of the corresponding formalism.

Formal power series of the following type ($m, n \in \mathbb{Z}$, $a_n, a_{m,n} \in A$),

$$a(z) = \sum_n \frac{a_n}{z^{n+1}}, \quad a(z, w) = \sum_{m,n} \frac{a_{m,n}}{w^{m+1} z^{n+1}}, \quad (123)$$

are called A -valued formal distributions in one, two, or more indeterminates $z, w, \dots$. We can freely multiply formal distributions by polynomials, but a product of two distributions in the same variable(s) is not generally defined. Also, there is a problem in treating negative powers of $(z - w)$ etc. as formal distributions. To do it unambiguously, we will use (when needed) the notation like

$$(z - w)_w^n \doteq \sum_{k \geq 0} (-)^k \binom{n}{k} w^k z^{n-k}, \quad (z - w)_z^n \doteq \sum_{k \geq 0} (-)^{n+k} \binom{n}{k} w^{n-k} z^k \quad (124)$$

which indicates, in positive powers of what variable (w or z in this case) the expansion is performed. So, e.g., $(z - w)_w^n$ is analogous to $(z - w)_{|z| > |w|}^n$ in terms of complex plane.

Of crucial importance in the present formalism is the δ -function:

$$\delta(z - w) = \delta(w - z) = \sum_n w^n z^{-n-1} = \dots + \frac{z}{w^2} + \frac{1}{w} + \frac{1}{z} + \frac{w}{z^2} + \dots \quad (125)$$

It exhibits standard-looking properties

$$f(z) \delta(z - w) = f(w) \delta(z - w), \quad (z - w) \delta(z - w) = 0, \quad \text{Res}_z f(z) \delta(z - w) = f(w) \quad (126)$$

where Res_z stands for a coefficient of z^{-1} , being an analog of $(2\pi i)^{-1} \oint dz$ on a complex plane. Note $\text{Res}_z \circ \partial_z = 0$. Evidently, the definition (125) mimics the well-known formulas like $\delta(x - y) = \sum_n e^{in(x-y)} = \sum_n (e^{ix})^n (e^{iy})^{-n}$.

Useful relations with the derivatives of the δ -function are listed below:

$$\frac{1}{n!} \partial_w^n \delta(z - w) = \frac{(-)^n}{n!} \partial_z^n \delta(z - w) = \sum_m \binom{m}{n} w^{m-n} z^{-m-1} \quad (n \geq 0) \quad (127)$$

$$(z - w)^m \partial_w^n \delta(z - w) = 0 \quad (m > n \geq 0) \quad (128)$$

$$(z - w)^m \frac{1}{n!} \partial_w^n \delta(z - w) = \frac{1}{(n-m)!} \partial_w^{n-m} \delta(z - w) \quad (0 \leq m \leq n) \quad (129)$$

$$\text{Res}_z f(z) \partial_w^n \delta(z - w) = \partial_w^n f(w), \quad \text{Res}_z \frac{(z - w)^m}{n!} \partial_w^n \delta(z - w) = \delta_{m,n} \quad (m, n \geq 0) \quad (130)$$

The following formula may also be of some use:

$$(z - w)_w^{-n} - (z - w)_z^{-n} = \frac{1}{(n-1)!} \partial_w^{n-1} \delta(z - w) \quad (n > 0). \quad (131)$$

Another principal notion here is locality. For the (most important) case of two variables the locality condition is

$$(z - w)^n a(z, w) = 0 \quad (n \geq N \geq 0). \quad (132)$$

It imposes severe restrictions on $a(z, w)$. Consider first the simplest case

$$\begin{aligned} (z - w) b(z, w) = 0 & \Rightarrow b_{m+1, n} = b_{m, n+1} \\ \Rightarrow b(z, w) = c(w) \delta(z - w), \quad c(w) = \text{Res}_z b(z, w). \end{aligned} \quad (133)$$

If now only $(z - w)^2 a(z, w) = 0$, we consecutively find that

$$\begin{aligned} (z - w) a(z, w) &= c_1(w) \delta(z - w) \equiv c_1(w) (z - w) \partial_w \delta(z - w), \\ c_1(w) &= \text{Res}_z (z - w) a(z, w), \\ a(z, w) - c_1(w) \partial_w \delta(z - w) &= c_0(w) \delta(z - w), \\ c_0(w) &= \text{Res}_z [a(z, w) - c_1(w) \partial_w \delta(z - w)] = \text{Res}_z a(z, w) \end{aligned}$$

and, therefore,

$$a(z, w) = c_0(w) \delta(z - w) + c_1(w) \partial_w \delta(z - w). \quad (134)$$

By induction, using (129) and (130), we show that $(z - w)^n a(z, w) = 0$ entails

$$a(z, w) = \sum_{m=0}^{n-1} c_m(w) \frac{1}{m!} \partial_w^m \delta(z - w), \quad c_m(w) = \text{Res}_z (z - w)^m a(z, w). \quad (135)$$

It is easily seen from (133) or (135) that a local formal distribution should contain infinitely many nonzero terms of positive and negative degree in both variables, otherwise it is identically zero. It is also clear that $a(w, z)$, $\partial_z a(z, w)$, $\partial_w a(z, w)$, $f(z) a(z, w)$ and $f(w) a(z, w)$ are local if $a(z, w)$ is.

One interesting example of a local distribution is provided by the product of two series with all unity coefficients:

$$\left(\sum_n z^{-n-1} \right) \left(\sum_m w^{-m-1} \right) = \delta(z - 1) \delta(w - 1) = \delta(z - w) \delta(w - 1) \quad (136)$$

(setting one argument of δ -function to a nonzero number makes sense).

One can verify (directly, or by induction) the following analog of the Taylor formula:

$$\left[f(z) - \sum_{n=0}^N \frac{(z - w)^n}{n!} \partial_w^n f(w) \right] \partial_w^N \delta(z - w) = 0. \quad (137)$$

It is precisely in this way we may consider an expression inside square brackets in (137) as something of the order $(z - w)^{N+1}$.

The following formulas with binomial coefficients are widely used in the calculations performed (or implicitly meant) in the present text:

$$\binom{-n-1}{m} = (-)^m \binom{n+m}{m} \quad (m \geq 0), \quad \binom{n}{m} = 0 \quad (m > n \geq 0), \quad (138)$$

$$\sum_{k=m}^n (-)^k \binom{n}{k} \binom{k}{m} = (-)^n \delta_{m,n}, \quad \sum_{k=0}^n (-)^k k^m \binom{n}{k} = (-)^n n! \delta_{m,n}, \quad (n \geq m \geq 0) \quad (139)$$

$$\sum_{k=0}^n (-)^k \binom{n}{k} \binom{p-k}{m} = \binom{p-n}{m-n}, \quad \sum_{k=0}^n (-)^k \binom{n}{k} \binom{p+k}{m} = (-)^n \binom{p}{m-n} \quad (140)$$

Operator product expansion

A single-variable formal distribution $a(z)$ is called a field on a vector space V if a_n are operators on V , and for any $v \in V$ a series $a(z)v$ contains only finitely many negative powers of z . In other words, $a_nv = 0$ for $n \geq N(v)$. Fields a and b are called (mutually) local if for some $N(a, b) \geq 0$

$$(z - w)^N [a(z), b(w)] = 0. \quad (141)$$

In what follows, we assume all fields to be pairwise local.

For each $n \in \mathbb{Z}$, an important notion of n -product of two fields a and b is introduced as follows:

$$(a_nb)(w) = \sum_m \frac{(a_nb)_m}{w^{m+1}}, \quad (a_nb)_m \doteq \sum_{k \geq 0} (-)^k \binom{n}{k} (a_{n-k}b_{m+k} - (-)^n b_{m+n-k}a_k). \quad (142)$$

It is seen from (142) that $(a_nb)(w)$ is also a local field, because for given n and v (or $c(z)$) we are able to find such N that $(a_nb)_mv = 0$ (resp. $(a_nb)_mc = 0$) for $m \geq N$.

Let us discuss the definition of the n -product in more details. For $n \geq 0$ it is equal to

$$(a_nb)(w) = \text{Res}_z (z - w)^n [a(z), b(w)]. \quad (143)$$

Locality condition (141) implies $(a_nb)(w) = 0$ ($n \geq N$) and, due to (135),

$$[a(z), b(w)] = \sum_{n=0}^{N-1} (a_nb)(w) \frac{1}{n!} \partial_w^n \delta(z - w). \quad (144)$$

Therefore, a commutator of local fields and (a finite number of) their n -products with non-negative n are closely related. This is also seen at the component level ($n \geq 0$):

$$(a_nb)_m = \sum_{k=0}^n (-)^k \binom{n}{k} [a_{n-k}, b_{m+k}], \quad [a_p, b_q] = \sum_{k \geq 0} \binom{p}{k} (a_kb)_{p+q-k} \quad (145)$$

(the second sum is also finite due to locality).

A special case $n = -1$ corresponds to the normal (or normally ordered) product,

$$(a_{-1}b)_m = \sum_{k < 0} a_k b_{m-k-1} + \sum_{k \geq 0} b_{m-k-1} a_k \Rightarrow (a_{-1}b)(w) = :a(w)b(w): \equiv :ab:(w), \quad (146)$$

where

$$:a(z)b(w): \doteq a_+(z)b(w) + b(w)a_-(z), \quad a_+(z) = \sum_{n < 0} \frac{a_n}{z^{n+1}}, \quad a_-(z) = \sum_{n \geq 0} \frac{a_n}{z^{n+1}}. \quad (147)$$

Usually $a_+(z)$ is said to contain creation and $a_-(z)$ annihilation operators. We can now verify that the following general formula for negative n agrees with the definition (142):

$$(a_nb)(w) = \frac{1}{(-n-1)!} : \partial^{-n-1} a(w) \cdot b(w) : \quad (n < 0) \quad (148)$$

There also exists a universal form of the n -product definition:

$$(a_nb)(w) = \text{Res}_z (a(z)b(w)(z - w)_w^n - b(w)a(z)(z - w)_z^n). \quad (149)$$

Remarkably, n -products play the role of coefficient functions of the operator product expansion (OPE). In CFT_2 , the latter is formulated for the radially ordered products

$$\mathcal{R} a(z) b(w) \doteq \begin{cases} a(z) b(w) & |z| > |w| \\ b(w) a(z) & |w| > |z| \end{cases} \quad (150)$$

Within the present formalism, OPE relations are expressed in terms of the definitions (124):

$$a(z) b(w) = \sum_n \frac{(a_n b)(w)}{(z-w)_w^{n+1}}, \quad b(w) a(z) = \sum_n \frac{(a_n b)(w)}{(z-w)_z^{n+1}} \quad (151)$$

or, equivalently,

$$a(z) b(w) = \sum_{n=0}^{N-1} \frac{(a_n b)(w)}{(z-w)_w^{n+1}} + :a(z) b(w): \quad (152)$$

$$b(w) a(z) = \sum_{n=0}^{N-1} \frac{(a_n b)(w)}{(z-w)_z^{n+1}} + :a(z) b(w): \quad (153)$$

$$:a(z) b(w): = \sum_{n \geq 0} (a_{-n-1} b)(w) (z-w)^n \quad (154)$$

Eqs. (154) and (151) are to be read not literally, but in the sense of Taylor's expansion (137), its coefficients given by (148). At the same time, (152) and (153) are derived from (131) and

$$a(z) b(w) - :a(z) b(w): = [a_-(z), b(w)], \quad (155)$$

$$b(w) a(z) - :a(z) b(w): = [b(w), a_+(z)] \quad (156)$$

by extracting negative and non-negative powers, respectively, from the z -expansion of (144), and are identically true. We see that a singular part of OPE is given by n -products with non-negative n , or by the commutator of fields.

It seems to be instructive to rewrite this derivation of OPE using the CFT_2 language. Let $\mathcal{R} a(z) b(w) = \sum_n \varphi_n(w) (z-w)^{-n-1}$. Then

$$\begin{aligned} \varphi_n(w) &= \text{Res}_{z=w} ((z-w)^n \mathcal{R} a(z) b(w)) \\ &= \frac{1}{2\pi i} \oint_w dz (z-w)^n \mathcal{R} a(z) b(w) = \frac{1}{2\pi i} \left(\oint_{|z|>|w|} dz - \oint_{|z|<|w|} dz \right) (z-w)^n \mathcal{R} a(z) b(w) \\ &= \frac{1}{2\pi i} \oint dz (a(z) b(w) (z-w)_{|z|>|w|}^n - b(w) a(z) (z-w)_{|z|<|w|}^n) \\ &= \text{Res}_z (a(z) b(w) (z-w)_{|z|>|w|}^n - b(w) a(z) (z-w)_{|z|<|w|}^n) = (a_n b)(w). \end{aligned} \quad (157)$$

It should be noted that from (142) and (145) (i.e., from the definition of the n -product plus the locality condition), the following useful formula (Borcherds identity) can be deduced. Namely, for $l, m, n \in \mathbb{Z}$, and a, b, c being fields, the following relations hold:

$$\sum_{k \geq 0} \binom{m}{k} (a_{l+k} b)_{m+n-k} c = \sum_{k \geq 0} (-)^k \binom{l}{k} (a_{m+l-k} (b_{n+k} c) - (-)^l b_{n+l-k} (a_{m+k} c)). \quad (158)$$

In CFT_2 , it's common practice to construct composite operators as normal products of two (or more) fields. To find then the OPE relations with these composite fields, the

so-called Wick formulas are used. In the present formalism the latter are obtained from (158), or (145), and look like

$$a_n(b_{-1}c) = b_{-1}(a_nc) + \sum_{k \geq 0} \binom{n}{k} (a_k b)_{n-k-1} c. \quad (159)$$

Vertex algebras

A vertex algebra consists of a vector space V (whose elements are called ‘states’), a vacuum state $\Omega \in V$, and a ‘state-field correspondence’ linear map Y which associates a field $a(z) \equiv Y(a, z) = \sum_n a_n z^{-n-1}$ (vertex operator) to each $a \in V$ in such a way that $Y(a_n b, z) = (a_n b)(z)$ are given by (142), all fields are mutually local ($a_n b = 0$ for $n \geq N(a, b)$), and $Y(\Omega, z)$ is equal to unity ($\Omega_n = \delta_{n,-1} \mathbf{1}$). In addition, a shift operator D is defined on V by $(Da)(z) \doteq \partial_z a(z)$. Note that an expression $a_n b$ is used here in two meanings: it denotes a state (an element in V obtained by the action of the operator a_n upon $b \in V$), and serves simultaneously as a name of the field $(a_n b)(z)$ (associated to this state via the map Y), which is exactly the n -product of two fields $a(z)$ and $b(z)$.

Now let us study the consequences of the above definitions. Due to locality of vertex operators, we may use not only (142), but also Borcherds’ relation (158). Evidently,

$$a_{-1} \Omega = a, \quad a_n \Omega = 0 \quad (n \geq 0), \quad D \Omega = 0, \quad (Da)_n = -n a_{n-1}. \quad (160)$$

Further, for $k \geq 0$,

$$(D^k a)_n = (-)^k k! \binom{n}{k} a_{n-k}, \quad a_{-k-1} = \frac{1}{k!} (D^k a)_{-1} \quad \Rightarrow \quad Y(a, z) \Omega = e^{zD} a. \quad (161)$$

Using (158) with $m = 0$, $n = -2$, $c = \Omega$, we obtain $D(a_l b) = (Da)_l b + a_l (Db)$, whence

$$[D, a_n] = (Da)_n = -n a_{n-1} \quad \Rightarrow \quad [D, Y(a, z)] = Y(Da, z) = \partial_z Y(a, z). \quad (162)$$

As an immediate generalization we find

$$e^{zD} Y(a, w) e^{-zD} = Y(e^{zD} a, w) = Y(a, w + z)_z. \quad (163)$$

For $m = -1$, $n = 0$, $c = \Omega$ formula (158) yields a so-called skew symmetry property

$$b_l a = \sum_{k \geq 0} \frac{(-)^{l+k+1}}{k!} D^k (a_{l+k} b) \quad \Rightarrow \quad Y(b, z) a = e^{zD} Y(a, -z) b. \quad (164)$$

The OPE relation assumes here the following form:

$$Y(a, z) Y(b, w) = Y(Y(a, z - w)_w b, w), \quad Y(b, w) Y(a, z) = Y(Y(a, z - w)_z b, w). \quad (165)$$

If $V = \bigoplus_{k \geq 0} V_k$ is a graded space equipped with a vertex algebra structure (so that $\Omega \in V_0$, $D : V_k \rightarrow V_{k+1}$, all a_n homogeneous), one usually rennumbers the components of the field $a(z)$ with $a \in V_\Delta$ in such a way (shifting index $n \rightarrow n - \Delta + 1$) that

$$a(z) = \sum_n a_n z^{-n-\Delta}, \quad a_{-n} : V_k \rightarrow V_{k+n}, \quad a_{-\Delta} \Omega = a, \quad a_n \Omega = 0 \quad (n > -\Delta). \quad (166)$$

As the famous example, the Virasoro algebra with central charge c can be reconstructed in this way from a single ‘conformal’ vector $\omega \in V_2$:

$$\omega(z) \equiv Y(\omega, z) = \sum_n \omega_n z^{-n-1} \equiv T(z) = \sum_n L_n z^{-n-2} \quad (\omega_n = L_{n-1}) , \quad (167)$$

$$L_{-2}\Omega = \omega , \quad L_{-1} = D , \quad L_0 a = \Delta a \quad (a \in V_\Delta) , \quad L_1 \omega = 0 , \quad L_2 \omega = \frac{c}{2} \Omega . \quad (168)$$

These definitions prove to be consistent, agree with (158) , and ultimately result in

$$[L_m, L_n] = (m-n)L_{m+n} + \frac{c}{12} m(m^2-1) \delta_{n,-m} , \quad (169)$$

$$\mathcal{R}T(z)T(w) = \frac{c}{2} (z-w)^{-4} + 2(z-w)^{-2}T(w) + (z-w)^{-1}\partial T(w) + \dots . \quad (170)$$

Free fermions

Below we describe the (Sugawara-like) construction of a conformal vector as a bilinear combination of ‘more elementary’ objects: in this (simplest) case, neutral fermions. The corresponding vertex algebra is generated by the vacuum state Ω and one more vector ϕ subject to

$$\phi_0\phi = \Omega , \quad \phi_n\phi = 0 \quad (n > 0) . \quad (171)$$

This exactly corresponds to

$$\phi(z) = \sum_n \phi_n z^{-n-1} , \quad [\phi_m, \phi_n]_+ = \delta_{m+n,-1} , \quad \phi(z)\phi(w) \sim \frac{1}{(z-w)_w} . \quad (172)$$

Since $\phi(z)$ is a fermionic field, we deal with anticommutators instead of commutators, and all formulas like (142), (145), and so on, should be modified accordingly.

Let us now introduce a vector

$$\omega \doteq \frac{1}{2} \phi_{-2}\phi \equiv \frac{1}{2} (D\phi)_{-1}\phi \equiv \frac{1}{2} \phi_{-2}\phi_{-1}\Omega \quad (173)$$

which is to be recognized as a conformal one. Eq. (159) yields

$$\phi_0\omega = -\frac{1}{2}D\phi , \quad \phi_1\omega = \frac{1}{2}\phi , \quad \phi_n\omega = 0 \quad (n > 1) \quad (174)$$

and (164) then gives

$$\omega_0\phi = D\phi , \quad \omega_1\phi = \frac{1}{2}\phi , \quad \omega_n\phi = 0 \quad (n > 1) \quad (175)$$

Analogously, we come to

$$\omega_0D\phi = D^2\phi , \quad \omega_1D\phi = \frac{3}{2}D\phi , \quad \omega_2D\phi = \phi , \quad (176)$$

and, finally, to

$$\omega_0\omega = D\omega , \quad \omega_1\omega = 2\omega , \quad \omega_3\omega = \frac{1}{4}\Omega \quad (177)$$

(only nonzero terms are displayed). So ω (173) is a genuine conformal vector, with central charge $c = 1/2$, and ϕ a *primary* element of conformal weight $\Delta = 1/2$ (that means literally (175)), with the OPE as follows:

$$\omega(z) \phi(w) = \frac{\phi(w)}{2(z-w)_w^2} + \frac{\partial\phi(w)}{(z-w)_w} + \dots, \quad (178)$$

$$\omega(z) \omega(w) = \frac{1}{4(z-w)_w^4} + \frac{2\omega(w)}{(z-w)_w^2} + \frac{\partial\omega(w)}{(z-w)_w} + \dots. \quad (179)$$

Note that in this case, unlike (166) and (167), we prefer not to shift indices of the field components in (172).

As for the vertex algebra as a whole, it is spanned, according to the Pauli principle, by the (finite) monomials of the form

$$\dots \phi_{-m}^{i_m} \dots \phi_{-1}^{i_1} \Omega \quad (m > 0, \quad i_m = 0, 1). \quad (180)$$

The second equality in (145) gives

$$[\omega_0, \phi_n] = (D\phi)_n = -n\phi_{n-1}, \quad [\omega_1, \phi_n] = \left(-\frac{1}{2} - n\right)\phi_n, \quad (181)$$

and one readily verifies that

$$\omega_0 a = Da, \quad \omega_1 a = \Delta a \quad (182)$$

for any a of the type (180), in accordance with the definition (168) of conformal vector.

Let us now proceed with a slightly more involved example related to charged fermions. Here the vertex algebra is generated by the vacuum state Ω and two fermionic (odd) states ψ^+ ($\equiv \psi$) and ψ^- ($\equiv \bar{\psi}$) subject to

$$\psi_0^+ \psi^- = \psi_0^- \psi^+ = \Omega, \quad \psi_0^+ \psi^+ = \psi_0^- \psi^- = \psi_n^\pm \psi^\pm = \psi_n^\pm \psi^\mp = 0 \quad (n > 0) \quad (183)$$

or, in other words (note renumbering ψ_n^+ w.r.t. (60)),

$$\psi^\pm(z) = \sum_n \psi_n^\pm z^{-n-1}, \quad [\psi_m^\pm, \psi_n^\pm]_+ = 0, \quad [\psi_m^\pm, \psi_n^\mp]_+ = \delta_{m+n,-1}, \quad (184)$$

$$\psi^\pm(z) \psi^\pm(w) \sim 0, \quad \psi^+(z) \psi^-(w) \sim \psi^-(z) \psi^+(w) \sim \frac{1}{(z-w)_w}. \quad (185)$$

Now we can construct a bilinear (bosonic) current (67)

$$J = \psi_{-1}^+ \psi^- = \psi_{-1}^+ \psi_{-1}^- \Omega \quad (186)$$

and, for arbitrary complex number λ , a conformal vector

$$\omega_\lambda = \frac{1}{2} J_{-1} J + \left(\frac{1}{2} - \lambda\right) D J = (1 - \lambda) \psi_{-2}^+ \psi^- + \lambda \psi_{-2}^- \psi^+ = [(1 - \lambda) \psi_{-2}^+ \psi_{-1}^- + \lambda \psi_{-2}^- \psi_{-1}^+] \Omega. \quad (187)$$

We will see that ψ^+ and ψ^- are primary states of conformal weights λ and $1 - \lambda$, respectively, whereas J has $\Delta = 1$ and becomes primary only for $\lambda = 1/2$ (when $c = 1$).

First, we check that two definitions of ω_λ (in terms of J and $\psi^\pm$) do agree. Using (142), (184), and properties of D , we find

$$J_{-1} J = (\psi_{-1}^+ \psi^-)_{-1} J = \sum_{k \geq 0} (\psi_{-1-k}^+ \psi_{-1+k}^- - \psi_{-2-k}^- \psi_k^+) \psi_{-1}^+ \psi_{-1}^- \Omega = (\psi_{-2}^+ \psi_{-1}^- + \psi_{-2}^- \psi_{-1}^+) \Omega,$$

$$DJ = (D\psi^+)_{-1}\psi^- + \psi_{-1}^+ D\psi^- = (\psi_{-2}^+\psi_{-1}^- + \psi_{-1}^+\psi_{-2}^-)\Omega = (\psi_{-2}^+\psi_{-1}^- - \psi_{-2}^-\psi_{-1}^+)\Omega,$$

which verifies (187).

Now we are ready to obtain, quite straightforwardly, the n -products (with $n \geq 0$), (anti)commutators, and singular parts of the OPE for $\psi^\pm$, J and ω_λ in all the relevant combinations (as usual, only nonzero contributions are listed):

$$\psi_0^\pm J = \mp \psi^\pm, \quad J_0 \psi^\pm = \pm \psi^\pm, \quad [J_m, \psi_n^\pm] = \pm \psi_{m+n}^\pm, \quad (188)$$

$$\psi^\pm(z)J(w) \sim \mp \frac{\psi^\pm(w)}{(z-w)_w}, \quad J(z)\psi^\pm(w) \sim \pm \frac{\psi^\pm(w)}{(z-w)_w}, \quad (189)$$

$$J_1 J = \Omega, \quad [J_m, J_n] = m\delta_{m,-n}, \quad J(z)J(w) \sim \frac{1}{(z-w)_w^2}, \quad (190)$$

$$\omega_{\lambda 0} \psi^\pm = D\psi^\pm, \quad \omega_{\lambda 1} \psi^+ = \lambda \psi^+, \quad \omega_{\lambda 1} \psi^- = (1-\lambda) \psi^-, \quad (191)$$

$$\omega_\lambda(z)\psi^\pm(w) \sim \frac{[1 \pm (2\lambda - 1)]\psi^\pm(w)}{2(z-w)_w^2} + \frac{\partial \psi^\pm(w)}{(z-w)_w}, \quad (192)$$

$$J_1 \omega_\lambda = J, \quad J_2 \omega_\lambda = (1-2\lambda)\Omega, \quad J(z)\omega_\lambda(w) \sim \frac{1-2\lambda}{(z-w)_w^3} + \frac{J(w)}{(z-w)_w^2}, \quad (193)$$

$$\omega_{\lambda 0} J = DJ, \quad \omega_{\lambda 1} J = J, \quad \omega_{\lambda 2} J = (2\lambda - 1)\Omega, \quad (194)$$

$$\omega_\lambda(z)J(w) \sim \frac{2\lambda - 1}{(z-w)_w^3} + \frac{J(w)}{(z-w)_w^2} + \frac{\partial J(w)}{(z-w)_w}, \quad (195)$$

$$\omega_{\lambda 0} \omega_\lambda = D\omega_\lambda, \quad \omega_{\lambda 1} \omega_\lambda = 2\omega_\lambda, \quad \omega_{\lambda 3} \omega_\lambda = (-6\lambda^2 + 6\lambda - 1)\Omega, \quad (196)$$

$$\omega_\lambda(z)\omega_\lambda(w) \sim \frac{-6\lambda^2 + 6\lambda - 1}{(z-w)_w^4} + \frac{2\omega_\lambda(w)}{(z-w)_w^2} + \frac{\partial \omega_\lambda(w)}{(z-w)_w}, \quad (197)$$

the central charge here being $c = -12\lambda^2 + 12\lambda - 2 = 1 - 3(2\lambda - 1)^2$.

Let us now consider an important set of primary states V^m (the same as $|m\rangle$ in (58)):

$$V^m = \psi_{-m}^+ \psi_{-m+1}^+ \dots \psi_{-1}^+ \Omega, \quad V^{-m} = \psi_{-m}^- \psi_{-m+1}^- \dots \psi_{-1}^- \Omega \quad (m \geq 0). \quad (198)$$

Of course, $V^0 = \Omega$, $V^{\pm 1} = \psi^\pm$. Using (184) and (188), one easily checks that

$$J_{n>0} V^m = 0, \quad J_0 V^m = m V^m, \quad m J_{-1} V^m = D V^m. \quad (199)$$

The last equality is verified by induction (say, for $m > 0$):

$$J_{-1} V^1 = J_{-1} \psi_{-1}^+ \Omega = \psi_{-2}^+ \Omega + \psi_{-1}^+ J = (D\psi^+)_{-1} \Omega + \psi_{-1}^+ \psi_{-1}^+ \psi^- = D V^1, \\ \psi_{-m-1}^+ J_{-1} V^m = 0, \quad J_{-1} V^{m+1} = \psi_{-m-2}^+ V^m = \frac{1}{m+1} [D, \psi_{-m-1}^+] V^m = \frac{D V^{m+1}}{m+1}.$$

Now, from

$$\omega_{\lambda n} = \frac{1}{2} \left(\sum_{k<0} J_k J_{n-k-1} + \sum_{k \geq 0} J_{n-k-1} J_k \right) + \left(\lambda - \frac{1}{2} \right) n J_{n-1} \quad (200)$$

and, in particular,

$$\omega_{\lambda 0} = \sum_{k \geq 0} J_{-k-1} J_k, \quad \omega_{\lambda 1} = \frac{1}{2} J_0 J_0 + \left(\lambda - \frac{1}{2} \right) J_0 + \sum_{k>0} J_{-k} J_k, \quad (201)$$

we see that

$$\omega_{\lambda 0} V^m = J_{-1} J_0 V^m = m J_{-1} V^m = D V^m, \quad \omega_{\lambda 1} V^m = \left(\frac{m^2}{2} + m\lambda - \frac{m}{2} \right) V^m. \quad (202)$$

These relations, together with $\omega_{\lambda n} V^m = 0$ for $n > 1$, characterize V^m as a primary state of conformal weight $(\frac{m^2}{2} + m\lambda - \frac{m}{2})$.

There exists a remarkable representation for the field $V^m(z)$ in terms of J_n and an additional invertible operator Y (previously introduced in (105); not to be confused with the state-field map itself!) defined by

$$Y \psi_n^+ = \psi_{n-1}^+ Y, \quad Y \psi_n^- = \psi_{n+1}^- Y, \quad Y \Omega = V^1 \implies Y V^m = V^{m+1}, \quad V^m = Y^m \Omega. \quad (203)$$

From

$$J_{n \neq 0} = \sum_k \psi_k^+ \psi_{n-k-1}^-, \quad J_0 = \sum_{k \geq 0} (\psi_{-k-1}^+ \psi_k^- - \psi_{-k-1}^- \psi_k^+) \quad (204)$$

one immediately concludes that

$$J_n Y = Y J_n \quad (n \neq 0), \quad [J_0, Y] = Y \implies [J_0, Y^m] = m Y^m. \quad (205)$$

A lengthy argument based on the Schur lemma shows that

$$V^m(z) = : e^{m\varphi(z)} : = Y^m \exp(-m \sum_{n < 0} \frac{J_n}{nz^n}) z^{mJ_0} \exp(-m \sum_{n > 0} \frac{J_n}{nz^n}) \quad (206)$$

(cf. (110), (112)), where

$$\varphi(z) = \ln Y + J_0 \ln z - \sum_{n \neq 0} \frac{J_n}{nz^n}, \quad \varphi'(z) = J(z). \quad (207)$$

We treat $\ln Y$ as a creation operator in accordance with $[J_0, \ln Y] = 1$. Differentiating $V^m(z)$ is in agreement with (199): $\partial_z V^m(z) = m : J(z) V^m(z) : \equiv m (J_{-1} V^m)(z)$. Another identity stemming from (206), $V^m(z) \Omega = e^{zD} V^m = \exp(-m \sum_{n < 0} \frac{J_n}{nz^n}) V^m$, is also valid due to the properties of Schur polynomials.

Virasoro algebra in KdV

The famous Korteweg-de Vries (KdV) equation

$$u_t = \frac{1}{4} (u_{xxx} + 6uu_x) \quad (208)$$

is a (first non-trivial) member of an integrable hierarchy

$$L_{t_n} = [L_+^{(2n-1)/2}, L] \quad (t_1 = x, \quad t_2 = t) \quad (209)$$

generated by the Lax operator

$$L = \partial^2 + u, \quad u = u(t, x), \quad \partial = \partial_x. \quad (210)$$

The KdV equation corresponds to $n = 2$, and so is governed by the M -operator

$$L_+^{3/2} = \partial^3 + \frac{3}{2} u \partial + \frac{3}{4} u_x = \partial^3 + \frac{3}{4} (u \partial + \partial \circ u). \quad (211)$$

The hierarchy (209) may be also seen as a reduction of the KP hierarchy which reduces its immense phase space to the (unique) square root of L , with a single unknown function u :

$$L^{1/2} = \partial + \frac{1}{2}u\partial^{-1} - \frac{1}{4}u_x\partial^{-2} + \frac{1}{8}(u_{xx} - u^2)\partial^{-3} + \mathcal{O}(\partial^{-4}). \quad (212)$$

The following *modified* KdV equation (mKdV) is closely related to (208):

$$q_t = \frac{1}{4}(q_{xxx} - 6q^2q_x). \quad (213)$$

Namely, it is obtained from (208) by the Miura transformation

$$u = q_x - q^2 \quad \Rightarrow \quad L = \partial^2 + u = (\partial - q)(\partial + q) \quad (214)$$

which sends (208) to

$$(\partial - 2q)(q_t - \frac{1}{4}q_{xxx} + \frac{3}{2}q^2q_x) = 0.$$

Let us now look at these integrable equations from the Hamiltonian standpoint. Here

$$f_t = \{H, f\}, \quad \{F, G\} \doteq \int dx dy \frac{\delta F}{\delta u(x)} \{u(x), u(y)\} \frac{\delta G}{\delta u(y)}, \quad (215)$$

$$F(u + \varepsilon v) \doteq F(u) + \varepsilon \int dx v(x) \frac{\delta F}{\delta u(x)} + \mathcal{O}(\varepsilon^2) \quad \Rightarrow \quad \frac{\delta F}{\delta u(x)} = \sum_{n=0}^{\infty} (-\partial)^n \frac{\partial F}{\partial u^{(n)}}, \quad (216)$$

and $u^{(n)} \equiv \partial^n u$. Assuming locality of the Poisson bracket,

$$\{u(x), u(y)\} = -\Lambda(x) \delta(x - y), \quad (217)$$

we come to

$$\{G, F\} = \int dx \frac{\delta F}{\delta u(x)} \Lambda(x) \frac{\delta G}{\delta u(x)}, \quad u_t = \Lambda \frac{\delta H}{\delta u}. \quad (218)$$

Hamiltonians H_m of the KdV hierarchy are known to be of the form $\sim \int dx \text{Res} L^{m+\frac{1}{2}}$ ($\text{Res} = \text{Res}_\partial$ is a coefficient of ∂^{-1}). For example,

$$H_0 = \int dx u, \quad H_1 = \frac{1}{4} \int dx u^2, \quad H_2 = \frac{1}{16} \int dx (2u^3 - u_x^2). \quad (219)$$

Moreover, KdV is a bi-Hamiltonian system, possessing (at least) two local Poisson structures of the type (217):

$$\Lambda_1 = 2\partial, \quad \Lambda_2 = \frac{1}{2}\partial^3 + 2u\partial + u_x = \frac{1}{2}\partial^3 + u\partial + \partial \circ u. \quad (220)$$

Acting on adjacent Hamiltonians, these two Λ operators produce the same result:

$$u_x \equiv u_{t_1} = \Lambda_1 \frac{\delta H_1}{\delta u} = \Lambda_2 \frac{\delta H_0}{\delta u} = u_x \quad (\text{identity}) \quad (221)$$

$$u_t \equiv u_{t_2} = \Lambda_1 \frac{\delta H_2}{\delta u} = \Lambda_2 \frac{\delta H_1}{\delta u} = \frac{1}{4}(u_{xxx} + 6uu_x) \quad (\text{KdV}) \quad (222)$$

$$u_{t_m} = \Lambda_1 \frac{\delta H_m}{\delta u} = \Lambda_2 \frac{\delta H_{m-1}}{\delta u} = \text{m-th member of KdV hierarchy} \quad (223)$$

It is the (symmetry) properties of the second Poisson bracket provided by Λ_2 that appear to be the main subject of this section. To begin with, we use the Miura transformation (214) to discover that

$$\Lambda_2 = \frac{1}{2}\partial^3 + 2u\partial + u_x = (\partial - 2q)\left(-\frac{\partial}{2}\right)(-\partial - 2q). \quad (224)$$

Then, from (218) and

$$\frac{\delta u(x)}{\delta q(y)} = (\partial_x - 2q)\delta(x - y), \quad \frac{\delta G}{\delta q(x)} = \int dy \frac{\delta u(y)}{\delta q(x)} \frac{\delta G}{\delta u(y)} = (-\partial_x - 2q) \frac{\delta G}{\delta u(x)}$$

we see that $-\frac{1}{2}\partial$ is nothing but a Poisson structure of mKdV:

$$\{q(x), q(y)\} = \frac{1}{2}\partial_x \delta(x - y). \quad (225)$$

Indeed, the second KdV bracket immediately follows from (225), (214) and (129). It is also readily verified that

$$q_t = -\frac{1}{2}\partial \frac{\delta H_1(u(q))}{\delta q} = \frac{1}{4}(q_{xxx} - 6q^2 q_x). \quad (226)$$

Our next observation is a striking similarity between the Poisson brackets of (m)KdV and appropriate field commutators in CFT_2 . Let us rewrite the second KdV bracket as

$$\{u(x), u(y)\} = -\left(\frac{c}{12}\partial_x^3 + 2u(x)\partial_x + u_x(x)\right)\delta(x - y), \quad c = 6. \quad (227)$$

Compare (227) and (225) with local commutators of the energy-momentum tensor T and free bosonic field a (the same as J (190)), respectively:

$$[T(z), T(w)] = -\left(\frac{c}{12}\partial_z^3 + 2T(z)\partial_z + T_z(z)\right)\delta(z - w), \quad (228)$$

$$[a(z), a(w)] = -\partial_z \delta(z - w). \quad (229)$$

Now it looks only natural to mimic the corresponding CFT_2 definitions as follows:

$$u(x) = \sum_n x^{-n-2} L_n, \quad q(x) = \sum_n x^{-n-1} a_n, \quad (230)$$

to reproduce the Virasoro and Heisenberg algebras, up to inessential overall factors:

$$\{L_m, L_n\} = (m - n)L_{m+n} + \frac{c}{12}m(m^2 - 1)\delta_{n, -m}, \quad (231)$$

$$\{a_m, a_n\} = -\frac{m}{2}\delta_{n, -m}. \quad (232)$$

To recall the techniques, we show how to deduce (225) from (232) and (127):

$$\begin{aligned} \{q(x), q(y)\} &= \sum_{mn} x^{-m-1} y^{-n-1} \{a_m, a_n\} = -\frac{1}{2} \sum_m m x^{-m-1} y^{m-1} \\ &= -\frac{1}{2} \partial_y \delta(y - x) = \frac{1}{2} \partial_x \delta(x - y). \end{aligned} \quad (233)$$

By the way, we have shown that the Miura map (214) is a kind of Sugawara construction: it builds the stress-tensor-like object $u(x)$ out of the “free scalar field” $q(x)$.

The first link between the KdV Poisson bracket (227) and the Virasoro algebra is thus established. Let us try to elaborate it further. Recall that the Virasoro algebra (Vir)

$$[L_m, L_n] = (m - n)L_{m+n} + \frac{c}{12} m(m^2 - 1) \delta_{n, -m} \quad (234)$$

can be realized as a centrally extended algebra of vector fields $f(z) \partial_z$ on a circle,

$$[f \partial, g \partial] \doteq (f g' - f' g) \partial + \frac{c}{12} \text{Res}_z (f g'''), \quad f' \equiv \partial f, \quad (235)$$

with $L_m = z^{-m+1} \partial_z$. Here Res_z (coefficient of z^{-1}) may be thought of as $\oint \frac{dz}{2\pi i}$, which we will denote simply by $\int dz$. Note the property $\text{Res}_z f' = 0$. Now let us show that in terms of structure constants of the algebra (235), the bracket (227) assumes the form of Lie-Poisson (Kirillov) bracket $\{\lambda_i, \lambda_j\} = c_{ij}^k \lambda_k$. Viewing (235) as an expression for a component of the commutator in a Lie algebra, like $[p, q]^k = c_{ij}^k p^i q^j$, we may cast it into functional form, reserving a single discrete index 0 for central terms:

$$[f, g](x) = \int dy dz C(x, y, z) f(y) g(z), \quad [f, g]_0 = \int dy dz C_0(y, z) f(y) g(z), \quad (236)$$

whence

$$C(x, y, z) = \delta(x - z) \partial_y \delta(y - z) - \delta(x - y) \partial_z \delta(y - z), \quad C_0(y, z) = -\frac{c}{12} \partial_z^3 \delta(y - z). \quad (237)$$

Now a functional analog of the Lie-Poisson bracket arises:

$$\{\lambda(y), \lambda(z)\} = \lambda_0 C_0(y, z) + \int dx C(x, y, z) \lambda(x) = \left(\frac{\lambda_0 c}{12} \partial_y^3 + (\lambda(y) + \lambda(z)) \partial_y \right) \delta(y - z). \quad (238)$$

We assume $\lambda_0 = 1$, and observe that the bracket (227) may also be written as

$$\{u(x), u(y)\} = -\left(\frac{c}{12} \partial_x^3 + (u(x) + u(y)) \partial_x \right) \delta(x - y) \quad (239)$$

owing to

$$u(y) \partial_x \delta(x - y) = u(x) \partial_x \delta(x - y) + (y - x) u_x \partial_x \delta(x - y) = (u(x) \partial_x + u_x(x)) \delta(x - y). \quad (240)$$

Therefore, it is of the Lie-Poisson form.

Lie-Poisson brackets are closely connected with the coadjoint action of a Lie algebra on its dual. Let us now find a counterpart of the Poisson structure Λ_2 (220) in Vir^* , a dual space to Vir. Proceeding a bit more accurately than above, we denote a general element of the latter by a pair $(f, a) \equiv f \partial + ca$, with a being a number, and rewrite (235) as

$$[(f, a), (g, b)] = (f g' - f' g, \frac{1}{12} \text{Res}_z f g'''). \quad (241)$$

Let us also denote the adjoint action of an infinitesimal element (ε, ω) (that is, a commutator $[(\varepsilon, \omega), \cdot]$) simply by δ_ε . Then

$$\text{ad}_{(\varepsilon, \omega)}(f, a) = (\delta_\varepsilon f, \delta_\varepsilon a), \quad \delta_\varepsilon f = \varepsilon f' - f \varepsilon' = (-f \partial + f') \varepsilon, \quad \delta_\varepsilon a = \frac{1}{12} \text{Res}_z f''' \varepsilon. \quad (242)$$

Further, let $(\lambda, \alpha) \in \text{Vir}^*$, α be a number, and the contraction be chosen in the form

$$\langle (f, a), (\lambda, \alpha) \rangle \doteq a\alpha + \text{Res}_z f\lambda. \quad (243)$$

The invariance condition for (243) is

$$\alpha \delta_\varepsilon a + a \delta_\varepsilon \alpha + \text{Res}_z (\lambda \delta_\varepsilon f + f \delta_\varepsilon \lambda) = 0. \quad (244)$$

This implies the following form of the coadjoint action $\delta_\varepsilon(\lambda, \alpha) \equiv \text{ad}_{(\varepsilon, \omega)}^*(\lambda, \alpha)$:

$$\delta_\varepsilon \alpha = 0, \quad \delta_\varepsilon \lambda = \left(\frac{\alpha}{12} \partial^3 + 2\lambda \partial + \lambda' \right) \varepsilon. \quad (245)$$

Thus, Λ_2 (220) corresponds (for $\alpha = 6$) to coadjoint action in Vir^* . This is in agreement with our previous observation that the bracket (227) is Lie-Poisson.

Moreover, eq. (245) provides additional evidence of an intimate relation between the KdV solutions $u(x)$ and the energy-momentum tensor $T(z)$ in CFT_2 . Recall the transformation rule of a primary field $\varphi(z)$ of conformal dimension Δ :

$$\tilde{\varphi}(\tilde{z})(d\tilde{z})^\Delta = \varphi(z)(dz)^\Delta, \quad \tilde{z} = z + \varepsilon(z). \quad (246)$$

For infinitesimal ε , this reads

$$\tilde{\varphi}(z) - \varphi(z) \simeq -(\Delta \varphi \partial + \varphi') \varepsilon. \quad (247)$$

Analogous formulas for the stress tensor (which corresponds to $\Delta = 2$ but is not a primary field due to an anomaly) look as follows:

$$\tilde{T}(\tilde{z})(d\tilde{z})^2 = T(z)(dz)^2 - \frac{c}{12}(dz)^2 \{\tilde{z}; z\} \quad (248)$$

with $\{\tilde{z}; z\}$ being the Schwarzian derivative

$$\{f(z); z\} \doteq \frac{f'''}{f'} - \frac{3}{2} \left(\frac{f''}{f'} \right)^2 \quad (249)$$

with the properties

$$\{t; z\} (dz)^2 = -\{z; t\} (dt)^2 = \{t; w\} (dw)^2 + \{w; z\} (dz)^2. \quad (250)$$

In the infinitesimal form,

$$\tilde{T}(z) - T(z) \simeq -\left(\frac{c}{12} \partial^3 + 2T\partial + T' \right) \varepsilon. \quad (251)$$

Comparing $\delta_\varepsilon f$ in (242) with (247), and $\delta_\varepsilon \lambda$ in (245) with (251), we see that vector fields (elements of Vir) belong to $\Delta = -1$ whereas their duals to $\Delta = 2$. Intrinsic duality of Δ and $1 - \Delta$ spaces is well known in CFT_2 . It is due to

$$\int dz f(z) \varphi(z) = \int f(z) (dz)^{1-\Delta} \varphi(z) (dz)^\Delta \quad (252)$$

being a natural invariant contraction.

All the reasoning given above suggests that the KdV solutions $u(x)$ also live in Vir^* and so belong to the class $\Delta = 2$ (with the central charge $c = 6$). Probably, the most direct

evidence of this fact is provided by the following observation. The Lax operator (210) reveals the covariant properties generalizing (246) with $\Delta = 2$,

$$\tilde{L}(\tilde{x}) = \partial_{\tilde{x}}^2 + \tilde{u}(\tilde{x}) = (\tilde{x}')^{-\frac{3}{2}} (\partial_x^2 + u(x)) \circ (\tilde{x}')^{-\frac{1}{2}}, \quad \tilde{x}' \equiv \frac{d\tilde{x}}{dx}, \quad (253)$$

if u transforms abnormally, like a conformal stress tensor:

$$\tilde{u}(\tilde{x})(d\tilde{x})^2 = u(x)(dx)^2 - \frac{1}{2}(dx)^2 \{\tilde{x}; x\}. \quad (254)$$

To show this, one uses

$$\partial_{\tilde{x}}^2 = \partial_{\tilde{x}} \circ \partial_{\tilde{x}} = (\tilde{x}')^{-1} \partial_x \circ (\tilde{x}')^{-1} \partial_x = (\tilde{x}')^{-2} \partial_x^2 - \tilde{x}'' (\tilde{x}')^{-3} \partial_x, \quad (255)$$

$$\partial_x^2 \circ (\tilde{x}')^{-\frac{1}{2}} = -\frac{1}{2} (\tilde{x}')^{-\frac{1}{2}} \{\tilde{x}; x\} - \tilde{x}'' (\tilde{x}')^{-\frac{3}{2}} \partial_x + (\tilde{x}')^{-\frac{1}{2}} \partial_x^2. \quad (256)$$

We conclude that not only the Poisson structure Λ_2 , but also the Lax operator of the KdV hierarchy, as well as all its solutions, exhibit apparent features of conformal covariance, being thus related with certain representations of the Virasoro algebra.